Targetry R&D

Harold G. Kirk
Brookhaven National Laboratory

Muon Collaboration Meeting
Shelter Island
May 8-14, 2002

Outline

1. Targetry Issues
2. The Targets
3. The Target System
4. Summary and Plans

Targetry Challenges

Study 1 Requirements

- 1.5 MW 16 GeV proton driver
- Low-Z Carbon target

Study 2 Requirements

- 1 MW 24 GeV proton driver
- High-Z Mercury target

Future upgrades

- 4 MW proton driver
- Target: ?

Capture in 20 T SC Solenoid

- Magnetohydrodynamic behavior of target
- Target dispersal in 20 T field

Target Types



- Low-Z: Less peak energy dissipation
- Mid-Z: Stronger material
- High-Z: Higher pion yield

Pion Production

High-Z is best for maximal soft pion production



However: Low-Z is best for higher momentum pion production





Low-Z Targets: Carbon



- Less peak energy dissipation
- Stationary Target Possible
- ORNL sublimation studies- P. Spampinato
- Doubtful beyond 1.5 MW

E951 Carbon Rods Test



ATJ



Carbon-Carbon



- The strain amplitudes for the anisotropic carbon-carbon composite are a factor of ~10 less than for ATJ carbon.

Mid-Z Targets: Alloys

B. King: Use robust moving metal targets



- Many different alloys available
- ANSYS beam/metal calculations- N. Simos
- Iron based alloys- P. Thieberger



Maximum initial stress as % of yield stress for the 1 MW option.
10000
1000
100
10
1
% of Yield Stress
Iron
Inconel 718
Vascomax 350
Super Invar
1
2
3
4
5
6
7
8
9
Target Radius (mm)

Fig. 3



Maximum initial stress as % of yield stress for the 4 MW option.
10000
1000
100
10
1
% of Yield Stress
Iron
Inconel 718
Vascomax 350
Super Invar
1
2
3
4
5
6
7
8
9
Target Radius (mm)

Fig. 4

Super-Invar Linear Expansion



Target Velocities and Lengths

	Velocity m/s	Length m
1 MW Option		
Super Invar	7.5	175
Vascomax C-350	0.85	16.8
4 MW Option		
Super Invar	12.0	274
Vascomax C-350	3.0	34

Brookhaven Linear Isotope Production Irradiation

One week of BLIP irradiation is equivalent to 10 years for a Super-invar moving target in a 1 MW facility.



Target consist of:

- 14 Tapered Super-invar rods
- 14 Tapered Inconnel rods
- 16 1mm diameter Nickel wires

High-Z Targets: Mercury

Key properties

- Maximal soft-pion production
- High pion absorption
- High peak energy dissipation
- Could be extended to 4 MW and beyond

Key issues

- Jet dynamics in a high-field solenoid
- Nature of target disruption
- Achievment of near-laminar flow for a 20 m/s jet

E951 Mercury Jet





Key Results

- Proton beam intensities upto 4 TP
- Dispersal velocities $\approx$ 10 m/s
- Dispersal delay $\approx$ 40 μs

E951
Windows Test Setup



Window Material:

10 mil Aluminum

11 mil Havar

13 mil Titanium Alloy

1 mm Inconel

6 mm Inconel

P.Thieberger

E951 Proton Beam Spot

New spot size from Autoradiography





Results: RMS $\sigma_x = 0.93$ mm ; $\sigma_y = 0.28$ mm



The CERN Passive Trough



Exposures at the AGS and at CERN ISOLDE. Results reported here by Adrian Fabich



T= 300 μs 600 μs 900 μs 1200 μs

Dispersal velocity measured as a function of proton pulse intensity and pulse separation.

Magnetohydrodynamic Jet Behavior

Initial Tests of a Mercury Jet in a 13 T Magnetic Field at the (CERN/Grenoble High Magnetic Field Laboratory)

4 mm diameter Hg Jet



$v = 4.6$ m/s, $B = 0$ T



$v = 4.0$ m/s, $B = 13$ T

Recent results reported here by Adrian Fabich

Additional Jet Studies



- Jet character needs improvement.
- Performance of alternative high-Z liquids: Woodsmetal
- Repeat beam exposures with 16 TP

- Build new 2 m/s horizontal jets (Woodsmetal and Hg)
- Build 10 m/s vertical Hg jet



The Targetry System



The Targetry Concept

- Capture the low-energy soft pions
- Use high-Z target
- Capture in a 20 T solenoid field
- Conduct pions into a 1.25 T decay channel

R&D Issues

- Jet dynamics in a solenoidal field
- Target dispersal in a high magnetic field
- Component performance in an intense radiation environment

E951

20T Pulsed Solenoid



1. 5 MVA Pulsed Power Supply
2. 15 Metric Tonnes Coil Package
3. 80° K operation
4. Switch power from outer to inner coil
5. Cost $4.5M

E951

14.5T Pulsed Solenoid

Windings, Coil Form & Cryostat for Cryogenic Pulse Magnet for 5 T, 10 T & 14.5 T



1. 2.2 MVA Pulsed Power Supply
2. 3.6 Metric Tonnes Coil Package
3. 30° K operation
4. Cost $1.5M

Future Plans

- Work with the AGS to achieve a single bunch intensity of 16 TP deliverable to the A3 line.
  - Repeat Hg jet target
  - Explore other solid target candidates (Inconel, Invar, Carbon, etc.)
- Develop a 20 m/s Hg jet.
- Develop and build a 2 m/s Woodsmetal jet.
- Develop and build a high-field pulsed solenoid.



E951 Pulsed Solenoid

Task Profile